

=> d his

(FILE 'HOME' ENTERED AT 14:11:25 ON 09 JUN 2005)

FILE 'REGISTRY' ENTERED AT 14:11:34 ON 09 JUN 2005

L1 1 S 97-72-3/RN

FILE 'CAPLUS' ENTERED AT 14:12:00 ON 09 JUN 2005

L2 864 S L1

L3 204 S L2 AND "ACETIC ANHYDRIDE"



L4 19 S L3 AND "ISOBUTYRIC ACID"

=>

=> d bib abs 1-19

L4 ANSWER 1 OF 19 CAPLUS COPYRIGHT 2005 ACS on STN
AN 2005:84839 CAPLUS
DN 142:113736
TI Preparation of novel esters of tertiary alcohols
IN Beldowicz, Maria; Kakol, Barbara; Kulig-Adamiak, Anna; Dornowski, Tadeusz;
Obukowicz, Bozena; Kaminski, Jaroslaw; Lewicka, Lidia; Kazimierczak,
Jerzy
PA Instytut Chemii Przemyslowej im. Prof. Ignacego Moscickiego, Pol.
SO Pol., 7 pp.
CODEN: POXXA7
DT Patent
LA Polish
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
	-----	----	-----	-----	-----
PI	PL 183367	B1	20020628	PL 1996-314863	19960619
PRAI	PL 1996-314863		19960619		
GI					



AB The title compds. I [R1-R3 = alkyl, Ph, alkylphenyl; R4 = alkyl; with provisos], potentially useful as flavorants in cosmetic and other industries, were prepared in one-pot synthesis by reacting freshly prepared Grignard reagent R1MgX [X = halo] with ketone R2C(O)R3 followed by reaction of the intermediate II with carboxylic acid derivative R4C(O)Y [Y = halo, OC(O)R; R = alkyl]. Thus, reacting Mg with PhBr in THF followed by addition of di-Et ketone and subsequently Ac2O afforded 90% III. The prepared compds. I demonstrated characteristic flavors which were evaluated on 1 through 8 point flavor scale.

L4 ANSWER 2 OF 19 CAPLUS COPYRIGHT 2005 ACS on STN
AN 2004:875971 CAPLUS
DN 141:351760
TI Dehydration process for making isobutyric anhydride from
isobutyric acid and acetic anhydride
IN Paul, Jean-Michel; Busca, Patrick
PA Atofina, Fr.
SO Eur. Pat. Appl., 6 pp.
CODEN: EPXXDW
DT Patent
LA French
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
	-----	----	-----	-----	-----
PI	EP 1468980	A1	20041020	EP 2004-290802	20040325

R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,
 IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, PL, SK, HR
 FR 2853900 A1 20041022 FR 2003-4785 20030416
 US 2005014974 A1 20050120 US 2004-824618 20040415
 JP 2004315536 A2 20041111 JP 2004-121366 20040416
 PRAI FR 2003-4785 A 20030416

AB A dehydration process is presented for making isobutyric anhydride from
isobutyric acid and **acetic anhydride**
 with distillation of the acetic acid byproduct.

RE.CNT 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 3 OF 19 CAPLUS COPYRIGHT 2005 ACS on STN
 AN 2004:673197 CAPLUS
 DN 141:182324
 TI Substituted clonidine derivatives. I. 3,5-Dichloro-4-(imidazolidin-2-ylideneamino)benzonitrile and 3,5-dichloro-4-(1,3-diisobutyrylimidazolidin-2-ylideneamino)benzonitrile
 AU Elssfah, E. M.; Chinnakali, K.; Fun, H. K.; Mathison, I. W.; Gan, E. K.; Zubaid, M.; Sam, T. W.; Khoo, K. S.
 CS X-ray Crystallography Unit, School of Physics, Universiti Sains Malaysia, Penang, 11800 USM, Malay.
 SO Acta Crystallographica, Section C: Crystal Structure Communications (1999), C55(7), iii, IUC9900066/1-2
 CODEN: ACSCEE; ISSN: 0108-2701
 URL: <http://journals.iucr.org/c/issues/1999/07/00/issconts.html>
 PB Munksgaard International Publishers Ltd.
 DT Journal
 LA English
 AB In the title compds., C10H8C12N4, (I), and C18H20C12N4O2, (II), the dihedral angle between the imidazolidine and Ph rings are 67.7(1) and 70.34(9)°, resp. In (I), the imidazolidine ring adopts a half-chair conformation, whereas in (II), it is in a flattened envelope conformation. In (I), the glide related mols. are linked by N2-H2A...N1(x, 1/2-y, z1/2) H bonds with an N...N distance of 2.848(4) Å to form an infinite chain along the c axis with the Ph rings stacked at a perpendicular distance of 3.633(5) Å. In (II), a weak C-H...N intramol. H bond exists between C11 and N1, with a C...N distance of 2.902(4) Å. Crystallog. data and atomic coordinates are given.

RE.CNT 10 THERE ARE 10 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 4 OF 19 CAPLUS COPYRIGHT 2005 ACS on STN
 AN 2002:657919 CAPLUS
 DN 137:195593
 TI Methods for the treatment of neuropathic pain by aryl nitron compounds
 IN Waterbury, David; Wood, Paul L.; Khan, M. Amin; Upasani, Ravindra B.
 PA Centaur Pharmaceuticals, Inc., USA
 SO PCT Int. Appl., 82 pp.
 CODEN: PIXXD2
 DT Patent
 LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2002065993	A2	20020829	WO 2002-US758	20020108
	WO 2002065993	A3	20021107		

W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN,
 CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
 GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
 LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH,

PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ,
UA, UG, US, UZ, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU,
TJ, TM

RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, CH,
CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR,
BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG

US 2002165274 A1 20021107 US 2002-43659 20020108

US 6835754 B2 20041228

US 2004209958 A1 20041021 US 2004-843298 20040512

PRAI US 2001-260469P P 20010108

US 2002-43659 A3 20020108

OS MARPAT 137:195593

AB Methods are disclosed for the treatment of neuropathic pain by aryl
nitron compounds. Method involves administration of an effective
neuropathic pain-treating dose of a pharmaceutical composition (Markush
structures are given). Substituted aryl nitron compounds are useful as
therapeutics for neuropathic pain conditions in mammals.

L4 ANSWER 5 OF 19 CAPLUS COPYRIGHT 2005 ACS on STN

AN 2002:169679 CAPLUS

DN 136:233214

TI Fiber-reinforced composite materials with good biodegradability

IN Takeishi, Hiromasa; Shibata, Mitsuhiro; Yosomiya, Tatsunori

PA Chiba Institute of Technology, Japan

SO Jpn. Kokai Tokkyo Koho, 9 pp.

CODEN: JKXXAF

DT Patent

LA Japanese

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	JP 2002069208	A2	20020308	JP 2000-258698	20000829
PRAI	JP 2000-258698		20000829		

AB The composite materials comprises surface-treated plant cellulose fibers
with OH group converted into esters and biodegradable thermoplastic
resins. Surface treating Manila hemp fibers with isobutyric anhydride and
pyridine, melt kneading 10 parts treated Manila hemp fibers with 90 parts
Biopol, and injection molding gave test pieces with flexural strength 26.3
MPa and flexural modulus 1460 MPa.

L4 ANSWER 6 OF 19 CAPLUS COPYRIGHT 2005 ACS on STN

AN 2001:167971 CAPLUS

DN 134:207727

TI Preparation of quinolinones and related bicyclic compounds as androgen and
progesterone receptor modulators.

IN Zhi, Lin; Tegley, Christopher; Pio, Barbara; Arjan van Oeveren, Cornelis;
Motamedi, Mehrnouch; Martinborough, Esther; West, Sarah; Higuchi, Robert;
Hamann, Lawrence; Farmer, Luc

PA Ligand Pharmaceuticals Incorporated, USA

SO PCT Int. Appl., 356 pp.

CODEN: PIXXD2

DT Patent

LA English


FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2001016108	A2	20010308	WO 2000-US23585	20000825
	WO 2001016108	A3	20011220		

W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN,
CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT,
LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU,
SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN,

YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY,
DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ,
CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG

US 6566372	B1	20030520	US 2000-649466	20000824
CA 2384435	AA	20010308	CA 2000-2384435	20000825
BR 2000013653	A	20020514	BR 2000-13653	20000825
EP 1212303	A2	20020612	EP 2000-959507	20000825
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL				
TR 200200508	T2	20020923	TR 2002-200200508	20000825
JP 2003508387	T2	20030304	JP 2001-519677	20000825
ZA 2002001053	A	20030528	ZA 2002-1053	20020206
NO 2002000912	A	20020429	NO 2002-912	20020225
BG 106539	A	20021031	BG 2002-106539	20020321
US 2003130505	A1	20030710	US 2002-299909	20021118
PRAI US 1999-150987P	P	19990827		
US 2000-649466	A3	20000824		
WO 2000-US23585	W	20000825		
OS MARPAT 134:207727				
GI				



AB Title compds., e.g. [I; R1, R2 = COR3, CSR3, SO2R3, NO, NR3R4, alkyl, alkenyl, haloalkyl, haloalkenyl, haloalkynyl, heteroalkyl, heteroalkenyl, heteroalkynyl, etc.; R1R2 = atoms to form (substituted) heterocyclyl; R3, R4 = H, (substituted) alkyl, alkenyl, alkynyl, haloalkyl, heteroalkyl, heteroaryl, aryl; R5 = H, F, Cl, Br, iodo, OR3, SR3, NR3R4, alkyl, haloalkyl, heteroalkyl; R6 = F, Cl, Br, iodo, Me, CF3, CHF2, cyano, CF2Cl, CF2OR3, OR3, SOR3, CO2R3, NR3R4, (substituted) alkyl, alkenyl, alkynyl, haloalkyl, heteroalkyl, etc.; R7, R8 = H, F, Cl, Br, iodo, cyano, OR3, NR3R4, SR3, SOR3, NR3COR4, alkyl, haloalkyl, heteroalkyl, etc.; R9 = H, F, Cl, iodo, OR3, NR3R4, SR3, SOR3, SO2R3, alkyl, haloalkyl, heteroalkyl; R10 = NR1R2, (substituted) heterocyclyl; Y = O, S, NR3, NOR3, CR3R4], were prepared. Thus, 6-amino-4-trifluoromethyl-2(1H)-quinolinone (preparation given) was stirred with propionaldehyde and NaBH3CN in MeOH to give 70-95% 6-propylamino-4-trifluoromethyl-2(1H)-quinolinone. The latter showed androgen receptor agonist activity with a potency of 27 nM. A drug composition is given.

L4 ANSWER 7 OF 19 CAPLUS COPYRIGHT 2005 ACS on STN
AN 1999:545441 CAPLUS
DN 131:207755
TI Material and method for manufacture of porous membrane for integration of large-scale integrated circuit
IN Aoi, Nobuo
PA Matsushita Electric Industrial Co., Ltd., Japan
SO Jpn. Kokai Tokkyo Koho, 12 pp.
CODEN: JKXXAF
DT Patent

LA Japanese

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
	-----	----	-----	-----	-----
PI	JP 11233506	A2	19990827	JP 1998-29431	19980212
	JP 3571522	B2	20040929		
	US 6194029	B1	20010227	US 1999-243491	19990203
	US 6319854	B1	20011120	US 2000-670698	20000928
PRAI	JP 1998-29431	A	19980212		
	US 1999-243491	A1	19990203		

AB The material for manufacture of the porous membrane comprises a solution containing

(A) silanol condensate particles and (B) (a) an alkyl-, halo-, OH-, or polar group (composed of C and heteroatom)-containing organic acid, (b) ≥ 2 OH-containing organic acids, (c) ≥ 1 OH-containing organic acid and ≥ 1 polar group-containing organic acid, or (d) an organic acid anhydride. The porous

membrane is manufactured by applying the above material on a substrate, followed by heating it. The organic acid catalyzes condensation of silanol condensate particles to improve the mech. strength of the membrane without blocking of the pores.

L4 ANSWER 8 OF 19 CAPLUS COPYRIGHT 2005 ACS on STN

AN 1999:460864 CAPLUS

DN 131:199342

TI Nonordinary destruction of aliphatic aldehydes C2-C4 in solutions of giant palladium clusters Pd-561

AU Gladii, S. L.; Starchevskii, M. K.; Lastoviyak, , Yu. V.; Pezderskii, Yu. A.; Vargaftik, M. N.; Moiseev, I. I.

CS Borislavsk. Naukovo-Dosl. Inst. "Sintez", Borislav, Ukraine

SO Dopovidi Natsional'noi Akademii Nauk Ukraini (1998), (1), 174-178

CODEN: DNAUFL; ISSN: 1025-6415

PB Prezidiya Natsional'noi Akademii Nauk Ukraini

DT Journal

LA Ukrain/Ukraine

AB Giant palladium cluster (Pd561) solns. are found to catalyze at 333 K and 0.1 MPa the oxidative destruction of aliphatic aldehydes C2-C4 yielding carbon dioxide and hydrocarbons. Acetaldehyde is converted to CO₂ and CH₄. Destruction of propanal, butanal and i-butanal yields CO₂ and olefins - accordingly, ethylene and propene. A reaction mechanism suggested includes the cleavage of the α -C-C bond of RCH₂-C=O coordinated with a Pd-atom.

L4 ANSWER 9 OF 19 CAPLUS COPYRIGHT 2005 ACS on STN

AN 1998:352133 CAPLUS

DN 129:54296

TI Preparation of N-acyl nitrogen-containing cyclic ketones

IN Ishikawa, Masahiro; Koike, Hitoshi

PA Yuki Gosei Kogyo Co., Ltd., Japan

SO Jpn. Kokai Tokkyo Koho, 7 pp.

CODEN: JKXXAF

DT Patent

LA Japanese

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
	-----	----	-----	-----	-----
PI	JP 10147571	A2	19980602	JP 1996-323615	19961119
PRAI	JP 1996-323615		19961119		

OS CASREACT 129:54296; MARPAT 129:54296

AB RCOA (A = 4- or 3-piperidone residue, 3-pyrrolidone residue; R = alkyl, aryl) are prepared by reaction of PhCH₂A (A = same as above) with RCO₂COR (R = same as above) in the presence of H donors or H and Pd catalysts. N-benzyl-4-piperidone was treated with Ac₂O in PhMe in the presence of

Pd/C under 5 kg/cm² H at 40° for 30 min to give 95%
N-acetyl-4-piperidone.

L4 ANSWER 10 OF 19 CAPLUS COPYRIGHT 2005 ACS on STN

AN 1995:980696 CAPLUS

DN 124:175094

TI On the potential of zeolites to catalyze aromatic acylation with
carboxylic acids

AU Gunnewegh, E. A.; Downing, R. S.; Van Bekkum, H.

CS Laboratory Organic Chemistry and Catalysis, Delft University Technology,
Delft, 2628 BL, Neth.

SO Studies in Surface Science and Catalysis (1995), 97(Zeolites: A Refined
Tool for Designing Catalytic Sites), 447-52

CODEN: SSCTDM; ISSN: 0167-2991

PB Elsevier

DT Journal

LA English

AB A symposium. The intramol. acylation of 4-phenylbutyric acid was
investigated as a model reaction for Friedel-Crafts acylation catalyzed by
zeolites in the liquid phase. This reaction is catalyzed by zeolite H-Beta
in 4-chlorotoluene as solvent. The catalytic ability of H-Beta was
demonstrated by the fact that at reflux temperature the total turnover number
(TON)

was found to be 35. However, the acylation of toluene or butylbenzene
with carboxylic acids was slow; the unbalanced adsorption equilibrium between
the two reactants on the H-Beta (Si/Al=12) may be a contributing factor in
this case. The acylation of anisole by carboxylic acids or acid
anhydrides however, was readily catalyzed by zeolite H-Beta.

L4 ANSWER 11 OF 19 CAPLUS COPYRIGHT 2005 ACS on STN

AN 1995:655207 CAPLUS

DN 123:56502

TI Method for manufacture of sucrose fatty acid esters with high degree of
substitution

IN Takamura, Yasuyuki; Murakami, Osamu

PA Dai Ichi Kogyo Seiyaku Co Ltd, Japan

SO Jpn. Kokai Tokkyo Koho, 6 pp.

CODEN: JKXXAF

DT Patent

LA Japanese

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
	-----	----	-----	-----	-----
PI	JP 07101974	A2	19950418	JP 1993-250258	19931006
PRAI	JP 1993-250258		19931006		

AB The title sucrose fatty acid esters are prepared by reaction of sucrose
and/or a sucrose fatty acid ester with a fatty acid anhydride, wherein the
reactants are dissolved in a solvent, made alkali, and reacted under an
inert gas. The preferred solvent is a basic organic solvent which is used at
≥1.0 mol/1.0 mol fatty acid anhydride. The preferred reaction
temperature is ≥30 to <120°. This process gives sucrose fatty
acid esters with high degree of substitution and markedly good hue. A
basic organic solvent makes the reaction system alkali, prevents coloration
due to the thermal decomposition of sucrose fatty acid esters, dissolves
sucrose and/or sucrose fatty acid esters to make the reaction system
homogeneous and improve the reaction rate, serves as an esterification
catalyst, and eliminates the need of using an alkali metal hydroxide or
carbonate, which also results in preventing coloration. Carrying out the
reaction under an inert gas prevents coloration due to the oxidation of
sucrose. Thus, sucrose 0.1, Ac₂O 1.6, and pyridine 4.8 mol were placed in
a flask and N was bubbled at 10 mL/min into the mixture with stirring to
make the O concentration 0.1 mg-O/L in the gas phase. The esterification was
carried out, while bubbling N into the mixture at 5 mL/min and the reaction

mixture was distilled in vacuo at ≤ 3 mmHg and 60° to remove pyridine, Ac₂O, and formed AcOH, and stirred with 200 g toluene and 300 g 0.5 weight% aqueous NaHCO₃ at 50° for 30 min followed by separating the toluene layer, washing it with H₂O, and distilling toluene at 60° at ≤ 3 mmHg to give sucrose acetate with 7.98 degree of substitution and APHA (American Public Health Association) color scale 200.

L4 ANSWER 12 OF 19 CAPLUS COPYRIGHT 2005 ACS on STN

AN 1995:490129 CAPLUS

DN 123:9071

TI Manufacture of carbamate group-containing esters

IN Jaawarudo, Efu Gurahe; Arutaa, Rakobitsutsu

PA Dainippon Ink & Chemicals, Japan

SO Jpn. Kokai Tokkyo Koho, 9 pp.

CODEN: JKXXAF

DT Patent

LA Japanese

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	JP 07017940	A2	19950120	JP 1993-164552	19930702
PRAI	JP 1993-164552		19930702		
OS	CASREACT 123:9071; MARPAT 123:9071				
AB	Treating carbamate-containing alcs. HOXO ₂ CNHR ₂ (R ₂ = H, C ₁ -18 organic group; X = C ₁ -12 difunctional group) with acid anhydrides (R ₁ CO) ₂ O (R ₁ = same as R ₂) in the presence of a base catalyst gives title esters R ₁ CO ₂ XO ₂ CNHR ₂ or treating HOXNHCO ₂ R ₂ with (R ₁ CO) ₂ O give R ₁ CO ₂ XNHCO ₂ R ₂ . Thus, treating hydroxyethylurethane with Ac ₂ O in the presence of NaOMe at $80-85^\circ$ gave .apprx.91.5% 2-carbamoyloxyethyl acetate.				

L4 ANSWER 13 OF 19 CAPLUS COPYRIGHT 2005 ACS on STN

AN 1994:455895 CAPLUS

DN 121:55895

TI O-derivatized alginic acid antigens

IN Pier, Gerald

PA Brigham and Women's Hospital, Inc., USA

SO PCT Int. Appl., 45 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 2

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 9408617	A1	19940428	WO 1993-US9909	19931015
	W: AU, CA, JP				
	RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
	AU 9454057	A1	19940509	AU 1994-54057	19931015
PRAI	US 1992-962480	A	19921016		
	WO 1993-US9909	W	19931015		
AB	An O-derivatized alginic acid antigen capable of eliciting opsonizing antibodies in vivo is disclosed. The O-derivatized antigen shows enhanced antigenicity and immunogenicity relative to native, non-O-derivatized alginic acid antigens, particularly the mucoid exopolysaccharide antigen of Pseudomonas aeruginosa. Pharmaceutical compns. containing the O-derivatized antigen of the invention are also described. The invention also pertains to use of the compns. as vaccines, in immunodiagnostic assays, and in methods for producing monoclonal antibodies reactive against the antigen. Methods for forming O-derivatized alginic acid antigens of the invention are described. A non-O-derivatized starting material is reacted in a solvent with an alkyl anhydride derived from a short chain fatty acid (e.g. propionic, isobutyric, or butyric acid).				

L4 ANSWER 14 OF 19 CAPLUS COPYRIGHT 2005 ACS on STN
AN 1993:602807 CAPLUS
DN 119:202807

TI Cobalt(II)-catalyzed reaction of aldehydes with **acetic anhydride** under an oxygen atmosphere: scope and mechanism
AU Bhatia, Beena; Punniyamurthy, T.; Iqbal, Javed
CS Dep. Chem., Indian Inst. Technol., Kanpur, 208016, India
SO Journal of Organic Chemistry (1993), 58(20), 5518-23
CODEN: JOCEAH; ISSN: 0022-3263

DT Journal

LA English

OS CASREACT 119:202807

AB The reaction of aldehydes with **acetic anhydride** in the presence of catalytic cobalt(II) chloride under an oxygen atmospheric at ambient temperature is dependent upon the reaction medium. Aliphatic aldehydes react in acetonitrile to give 1,2-diones whereas the aromatic aldehydes are acylated to yield the corresponding acylals. On the other hand, carboxylic acids are obtained from aliphatic and aromatic aldehydes by conducting the reaction in dichloromethane or benzene. Cobalt(II) chloride in acetonitrile catalyzes the conversion of aliphatic aldehydes to the corresponding anhydrides in the absence of **acetic anhydride** whereas aromatic aldehydes remain largely unaffected under these conditions. A preliminary mechanistic study in three different solvents (i.e. acetonitrile, dichloroethane, and DMF) has revealed that in acetonitrile and in the presence of **acetic anhydride**, aliphatic aldehydes behave differently than aromatic aldehydes. Some trapping expts. using Me acrylate and stilbene have been conducted to demonstrate the occurrence of an acyl cobalt and peroxyacyl cobalt intermediate during these reactions.

L4 ANSWER 15 OF 19 CAPLUS COPYRIGHT 2005 ACS on STN
AN 1992:614970 CAPLUS
DN 117:214970

TI Method for purification of carboxylic acids and anhydrides
IN Zoeller, Joseph Robert; Moncier, Regina Michelle
PA Eastman Kodak Co., USA
SO PCT Int. Appl., 13 pp.
CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 1



	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 9212954	A1	19920806	WO 1992-US631	19920127
	W: CA, JP, KR				
	RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LU, MC, NL, SE				
	US 5175363	A	19921229	US 1991-646029	19910128
	CA 2098293	AA	19920729	CA 1992-2098293	19920127
	EP 569492	A1	19931118	EP 1992-905057	19920127
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, MC, NL, SE				
	JP 06505257	T2	19940616	JP 1992-505713	19920127
PRAI	US 1991-646029	A	19910128		
	WO 1992-US631	W	19920127		

AB A method for reducing the amount of olefinic impurity in the title C2-8 carboxylic acids and C4-16 anhydrides comprises contacting them with a strong acidic resin. A sample of AcOH contaminated with 221 ppm 1-octene was added to Amberlyst-15, the mixture refluxed for 3 h to give AcOH containing only 15 ppm 1-octene.

L4 ANSWER 16 OF 19 CAPLUS COPYRIGHT 2005 ACS on STN
AN 1992:135528 CAPLUS

DN 116:135528
 TI Performance-oriented packaging standards; changes to classification, hazard communication, packaging and handling requirements based on UN standards and agency initiative
 CS United States Dept. of Transportation, Washington, DC, 20590-0001, USA
 SO Federal Register (1990), 55(246), 52402-729, 21 Dec 1990
 CODEN: FEREAC; ISSN: 0097-6326
 DT Journal
 LA English
 AB The hazardous materials regulations under the Federal Hazardous Materials Transportation Act are revised based on the United Nations recommendations on the transport of dangerous goods. The regulations cover the classification of materials, packaging requirements, and package marking, labeling, and shipping documentation, as well as transportation modes and handling, and incident reporting. Performance-oriented stds. are adopted for packaging for bulk and nonbulk transportation, and SI units of measurement generally replace US customary units. Hazardous material descriptions and proper shipping names are tabulated together with hazard class, identification nos., packing group, label required, special provisions, packaging authorizations, quantity limitations, and vessel stowage requirements.

L4 ANSWER 17 OF 19 CAPLUS COPYRIGHT 2005 ACS on STN
 AN 1988:406370 CAPLUS
 DN 109:6370
 TI Thermal transformations of some 3-acyl-2-oxo-2H-1-benzopyrans with acid anhydrides
 AU Bodzhilova, A.; Ivanov, Kh.
 CS Fac. Chem., Univ. Sofia, Sofia, 1126, Bulg.
 SO Indian Journal of Chemistry, Section B: Organic Chemistry Including Medicinal Chemistry (1987), 26B(8), 731-5
 CODEN: IJSBDB; ISSN: 0376-4699
 DT Journal
 LA English
 OS CASREACT 109:6370
 GI



AB The reaction of 3-benzoyl- and 3-acetyl-2-oxo-2H-1 benzopyrans I (R = Ph, Me) with acid anhydrides in the presence of AcONa or Et3N was studied. Thus, with propionic, butyric and **isobutyric acid** anhydrides, I (R = Ph) affords the rearrangement products II and III (R1 = H, R2 = Me, Et; R1 = R2 = Me), dilactones IV and benzyl ester V.

L4 ANSWER 18 OF 19 CAPLUS COPYRIGHT 2005 ACS on STN
 AN 1977:71270 CAPLUS
 DN 86:71270
 TI Kinetics and mechanism of the reaction of aliphatic nitriles with carboxylic acids
 AU Zil'berman, E. N.; Navolokina, R. A.; Minchuk, F. F.; Danov, S. M.; Gromova, G. V.
 CS Gor'k. Politekh. Inst. im. Zhdanova, Dzerzhinsk, USSR
 SO Izvestiya Vysshikh Uchebnykh Zavedenii, Khimiya i Khimicheskaya Tekhnologiya (1976), 19(9), 1395-8
 CODEN: IVUKAR; ISSN: 0579-2991
 DT Journal
 LA Russian
 AB Rate consts. and activation parameters were determined for the reaction of RCN with RCO2H (R = Me, Et, Pr, Me2CH) to give RCONH2 and (RCO)2O and for the further reaction of these products to give (RCO)2NH and RCO2H. A 4-center cyclic transition state was suggested for the 2nd step.

L4 ANSWER 19 OF 19 CAPLUS COPYRIGHT 2005 ACS on STN
 AN 1952:57248 CAPLUS
 DN 46:57248
 OREF 46:9565f-g
 TI Function of pyridine in the carboxylic acid-thionyl chloride system
 AU Gerrard, W.; Thrush, A. M.
 CS Northern Polytech., London
 SO Journal of the Chemical Society, Abstracts (1952) 741-2
 CODEN: JCSAAZ; ISSN: 0590-9791
 DT Journal
 LA Unavailable
 AB The addition of 0.5 mol. SOCl2 to 1 mol. AcOH and 1 mol. C5H5N in ether at -10° gives an immediate precipitate which appears to be an equimol. mixture of C5H5N.HCl and C5H5N.HOSOC1; the ether yields 97% Ac2O. In the same way (iso-PrCO)2O, (iso-BuCO)2O, and (C6H13CO)2O were obtained in 98, 99, and 97% yields. A postulated mechanism involves the 4-center broadside approach of the reacting mols.

=>